

DrugCheck®

Drug Cup Cutoff Levels/Detection Times

Drug Type	Pharmaceutical Names	Street Names	Urine Detection Times	Signs of Use
Amphetamine (AMP)	Dexedrine, Benzedrine, Dextrostat, Adderall, Biphphetamine	Speed, Bennies, Dexies, Uppers, Footballs	Cutoff level - 1000 ng/ml Detection Times: 2-4 days	Dilated pupils, decreased appetite, dry mouth & nose, Bad breath, Frequent lip licking, talkative, nervous Increase in heart rate & blood pressure
Methamphetamine (MET)	Desoxyn, Methedrine	Ice, Crystal, Meth, Doe	Cutoff level - 1000 ng/ml Detection Times: 2-4 days	Same as amphetamines Overdoses cause elevated body temp.
Ecstasy (MDMA)	None	XTC, E, Adam, Blue lips, Blue kisses Disco biscuits, X-ing, Doctor	Cutoff level - 500 ng/ml Detection Times: 1-3 days	Same as amphetamines May grind teeth and clench jaw, Sweating and increase in body temp.
MDA	None	Eve, Love Pill	Detection Times : 1-3 days	Same as amphetamines Hallucinations, increase in body temp. Tachycardia, agitation, very talkative
Cocaine (COC)	None	Coke, Crack, Snow, Flake, Rock, Nose Candy, Powder, Blow	Cutoff level - 300 ng/ml Detection Times: 2-4 days	Dilates pupils, increase heart rate & BP Anesthesia, euphoria, sensitive to sound & touch Less need for food, cardiotoxic
Marijuana (THC)	Marinol	Weed, Pot, Grass, Reefer, Hash, Mary Jane, Cannabis, Columbian Acapulco Gold, Ganja, Bhang	Cutoff level - 50 ng/ml Detection Times: 1-2 joints: 2-3 days Oral Ingestion: 1-5 days Moderate smoker: 5 days Heavy smoker (daily): 10 days Chronic smoker-5 joints per day 4-18 days or as high as a month	Increase appetite, altered perception, memory loss Bloodshot eyes, giggle frequently, trouble walking Disorientation, increase heart rate, vasodilatation Dry mouth & throat, decrease respiratory rate Panic attacks, & paranoia
Phencyclidine (PCP)	Sernylan	Veterinary anesthetic Angel dust, Hog, Killer Weed	Cutoff level - 25 ng/ml Detection Times: Up to 14 days Up to 30 days with chronic users	Dissociative anesthesia-analgesia & amnesia Dissociated from environment but not asleep Pulse & BP elevated, coma
Ketamine	Ketalar	Veterinary Anesthetic Kat, Special K, Vitamin K, K, Horse Tranquilizer	Cutoff level - 25 ng/ml Detection Times: Up to 14 days Up to 30 days with chronic users	Analgesia, amnesia, hallucinations, Foul-tasting drip from nasal passages Users feel they are outside of their body, vomiting Seizures, dissociation
Gamma Hydroxybutyric Acid, (GHB)	None	GHB, Cherry meth, Easy Lay, jib Georgia Home Boy, sleep 500, Soap, Vita G, Grievous Bodily Harm Scoop, Goop, Fantasy, Everclear Salty water, Liquid X	Detection Times: 1 day	Users feel euphoric then sleepy Used as a date rape drug Slows breathing and heart rate

Drug Type	Pharmaceutical Names	Street Names	Urine Detection Times	Signs of Use
Opiates (OPI 300) (OPI 2000)	Hydromorphone (Dilaudid), Codeine, Tylenol with Codeine, Morphine (Avinza Kadian, MS Contin, Oramorph SR) Hydrocodone (Vicodin) - Positive with 300 cutoff	(Heroin & Morphine based drugs) Smack, H, Skag, Dope, Junk, Horse	Cutoff level - 300 ng/ml or 2000 ng/ml Detection Times: 1-2 days	Pupils constricted, pulse & BP lowered Body temp. lowered, euphoria initially and then drowsy Respiratory depression, convulsions, constipation, coma
Oxycodone (OXY)	OxyContin, Percodan, Percocet Roxicodone, Tylox OxyContin - time released form of Oxycodone	OXY, OC,OX, Oxycotton, Hillbilly Heroin, Killer	Cutoff level - 100ng/ml Detection Times: 1-3 days OxyContin longer detection times	Same as Opiates
Propoxyphene (PPX)	Darvon, Darvocet	Yellow Footballs	Cutoff level - 300ng/ml Detection Times: 3-4 days	Same as Opiates
Barbiturates (BAR)	Nembutal, Luminal, Seconal, Tuinal, Florinal, Neodorm, Immenoctal, Stadodorm	Barbs, Downers, Tranqs, Blue Dolls, Bluebirds, Rainbows, Yellows, Reds, Red Devils, M&Ms	Cutoff level - 300 ng/ml Detection Times: Short acting - 1 day (Seconal, Nembutal) Long acting: 2-3 weeks (Phenobarb)	Pupils normal, but reaction to light slow Pulse, BP & respiratory rate lowered Sedation, stupor, coma
Methadone (MTD)	Dolophine, Methadone, L-Polamidon Physeptone	Amidone, Fizzes	Cutoff level - 300 ng/ml Detection Times: 2-3 days	Same as other Opiates Similar to heroin but less intense Sedation analgesia
Benzodiazepines (BZD)	Librium, Valium, Ativan, Halicon, Vivol, Novopoxide, Remestan, Rohypnol, Xanax, Tranxene, Restoril, Versed, Clonopin, Rivotril, Dalaman, Serax	Bennies, Rophies	Cutoff level - 300 ng/ml Detection Times: Normal dose - 2-3 days Chronic Use (1 or more yrs) 4-6 weeks	Drowsiness, dizziness, muscle relaxation Hangover effect after large dose Rohynol used as rape drug
Tricyclic Antidepressants (TCA)	Amitriptyline (Elavil, Endep), Cyclobenzaprine (Flexeril), Clomipramine (Anafranil), Perphenazine (Triafon), Desipramine (Norpramin), Imipramine (Totranil), Nortriptyline (Pamelor), Protriptyline (Vivactil), Trimipramine (Surmontil), Promazine (Sparine), Nordoxepine, Doxepin (Sinequan)		Cutoff level - 1000 ng/ml Detection Times: Therapeutic dose - 3 days Extended use - 4 weeks	Elevated body temperature, Elevated pulse & blood pressure. Pupils dilated, agitation, delirium, seizures Constipation, dry, flush skin

*Customer reference only. Not for medical use.